

Package ‘BivGeo’

January 20, 2025

Type Package

Title Basu-Dhar Bivariate Geometric Distribution

Version 2.0.1

Author Ricardo Puziol de Oliveira and Jorge Alberto Achcar

Maintainer Ricardo Puziol de Oliveira <rpuziol.oliveira@gmail.com>

Description Computes the joint probability mass function (pmf), the joint cumulative function (cdf), the joint survival function (sf), the correlation coefficient, the covariance, the cross-factorial moment and generate random deviates for the Basu-Dhar bivariate geometric distribution as well the joint probability mass, cumulative and survival function assuming the presence of a cure fraction given by the standard bivariate mixture cure fraction model. The package also computes the estimators based on the method of moments.

Depends R (>= 3.0.2)

Imports stats

URL For more details, see de Oliveira et. al (2018)
<doi:10.1285/i20705948v11n1p108>.

RoxygenNote 6.1.1

Encoding UTF-8

NeedsCompilation no

LazyData TRUE

License GPL (>= 2)

Repository CRAN

Date/Publication 2019-01-03 15:10:16 UTC

Contents

cfbivgeo	2
corbivgeo	3
covbivgeo	5
dbivgeo	6
dbivgeocure	8
mombivgeo	10

pbivgeo	12
pbivgeocure	14
rbivgeo	16

Index	19
--------------	-----------

cfbivgeo	<i>Cross-factorial Moment for the Basu-Dhar Bivariate Geometric Distribution</i>
----------	--

Description

This function computes the cross-factorial moment for the Basu-Dhar bivariate geometric distribution assuming arbitrary parameter values.

Usage

```
cfbivgeo(theta)
```

Arguments

theta vector (of length 3) containing values of the parameters θ_1, θ_2 and θ_3 of the Basu-Dhar bivariate Geometric distribution. For real data applications, use the maximum likelihood estimates or Bayesian estimates to get the cross-factorial moment.

Details

The cross-factorial moment between X and Y , assuming the Basu-Dhar bivariate geometric distribution, is given by,

$$E[XY] = \frac{1 - \theta_1\theta_2\theta_3^2}{(1 - \theta_1\theta_3)(1 - \theta_2\theta_3)(1 - \theta_1\theta_2\theta_3)}$$

Note that the cross-factorial moment is always positive.

Value

`cfbivgeo` computes the cross-factorial moment for the Basu-Dhar bivariate geometric distribution for arbitrary parameter values.

Invalid arguments will return an error message.

Author(s)

Ricardo P. Oliveira <rpuziol.oliveira@gmail.com>

Jorge Alberto Achcar <achcar@fmrp.usp.br>

Source

`cfbivgeo` is calculated directly from the definition.

References

Basu, A. P., & Dhar, S. K. (1995). Bivariate geometric distribution. *Journal of Applied Statistical Science*, **2**, 1, 33-44.

Li, J., & Dhar, S. K. (2013). Modeling with bivariate geometric distributions. *Communications in Statistics-Theory and Methods*, **42**, **2**, 252-266.

Achcar, J. A., Davarzani, N., & Souza, R. M. (2016). Basu–Dhar bivariate geometric distribution in the presence of covariates and censored data: a Bayesian approach. *Journal of Applied Statistics*, **43**, 9, 1636-1648.

de Oliveira, R. P., & Achcar, J. A. (2018). Basu-Dhar's bivariate geometric distribution in presence of censored data and covariates: some computational aspects. *Electronic Journal of Applied Statistical Analysis*, **11**, 1, 108-136.

de Oliveira, R. P., Achcar, J. A., Peralta, D., & Mazucheli, J. (2018). Discrete and continuous bivariate lifetime models in presence of cure rate: a comparative study under Bayesian approach. *Journal of Applied Statistics*, 1-19.

Examples

```
cfbivgeo(theta = c(0.5, 0.5, 0.7))
# [1] 2.517483
cfbivgeo(theta = c(0.2, 0.5, 0.7))
# [1] 1.829303
cfbivgeo(theta = c(0.8, 0.9, 0.1))
# [1] 1.277864
cfbivgeo(theta = c(0.9, 0.9, 0.9))
# [1] 35.15246
```

corbivgeo

Correlation Coefficient for the Basu-Dhar Bivariate Geometric Distribution

Description

This function computes the correlation coefficient analogous of the Pearson correlation coefficient for the Basu-Dhar bivariate geometric distribution assuming arbitrary parameter values.

Usage

```
corbivgeo(theta)
```

Arguments

theta vector (of length 3) containing values of the parameters θ_1, θ_2 and θ_3 of the Basu-Dhar bivariate Geometric distribution. For real data applications, use the maximum likelihood estimates or Bayesian estimates to get the correlation coefficient.

Details

The correlation coefficient between X and Y, assuming the Basu-Dhar bivariate geometric distribution, is given by,

$$\rho = \frac{(1 - \theta_3)(\theta_1\theta_2)^{1/2}}{1 - \theta_1\theta_2\theta_3}$$

Note that the correlation coefficient is always positive which implies that the Basu-Dhar bivariate geometric distribution is useful for bivariate lifetimes with positive correlation.

Value

[corbivgeo](#) computes the correlation coefficient analogous to the Pearson correlation coefficient for the Basu-Dhar bivariate geometric distribution for arbitrary parameter values.

Invalid arguments will return an error message.

Author(s)

Ricardo P. Oliveira <rpuziol.oliveira@gmail.com>

Jorge Alberto Achcar <achcar@fmrp.usp.br>

Source

[corbivgeo](#) is calculated directly from the definition.

References

- Basu, A. P., & Dhar, S. K. (1995). Bivariate geometric distribution. *Journal of Applied Statistical Science*, **2**, 1, 33-44.
- Li, J., & Dhar, S. K. (2013). Modeling with bivariate geometric distributions. *Communications in Statistics-Theory and Methods*, **42**, **2**, 252-266.
- Achcar, J. A., Davarzani, N., & Souza, R. M. (2016). Basu–Dhar bivariate geometric distribution in the presence of covariates and censored data: a Bayesian approach. *Journal of Applied Statistics*, **43**, 9, 1636-1648.
- de Oliveira, R. P., & Achcar, J. A. (2018). Basu-Dhar’s bivariate geometric distribution in presence of censored data and covariates: some computational aspects. *Electronic Journal of Applied Statistical Analysis*, **11**, 1, 108-136.
- de Oliveira, R. P., Achcar, J. A., Peralta, D., & Mazucheli, J. (2018). Discrete and continuous bivariate lifetime models in presence of cure rate: a comparative study under Bayesian approach. *Journal of Applied Statistics*, 1-19.

Examples

```
corbivgeo(theta = c(0.5, 0.5, 0.7))
# [1] 0.1818182
corbivgeo(theta = c(0.2, 0.5, 0.7))
# [1] 0.102009
corbivgeo(theta = c(0.8, 0.9, 0.1))
# [1] 0.822926
corbivgeo(theta = c(0.9, 0.9, 0.9))
# [1] 0.3321033
```

`covbivgeo`*Covariance for the Basu-Dhar Bivariate Geometric Distribution*

Description

This function computes the covariance for the Basu-Dhar bivariate geometric distribution assuming arbitrary parameter values.

Usage

```
covbivgeo(theta)
```

Arguments

`theta` vector (of length 3) containing values of the parameters θ_1, θ_2 and θ_3 of the Basu-Dhar bivariate Geometric distribution. For real data applications, use the maximum likelihood estimates or Bayesian estimates to get the covariance.

Details

The covariance between X and Y , assuming the Basu-Dhar bivariate geometric distribution, is given by,

$$Cov(X, Y) = \frac{\theta_1 \theta_2 \theta_3 (1 - \theta_3)}{(1 - \theta_1 \theta_3)(1 - \theta_2 \theta_3)(1 - \theta_1 \theta_2 \theta_3)}$$

Note that the covariance is always positive.

Value

`covbivgeo` computes the covariance for the Basu-Dhar bivariate geometric distribution for arbitrary parameter values.

Invalid arguments will return an error message.

Author(s)

Ricardo P. Oliveira <rpuziol.oliveira@gmail.com>

Jorge Alberto Achcar <achcar@fmrp.usp.br>

Source

`covbivgeo` is calculated directly from the definition.

References

Basu, A. P., & Dhar, S. K. (1995). Bivariate geometric distribution. *Journal of Applied Statistical Science*, **2**, 1, 33-44.

Li, J., & Dhar, S. K. (2013). Modeling with bivariate geometric distributions. *Communications in Statistics-Theory and Methods*, **42**, **2**, 252-266.

Achcar, J. A., Davarzani, N., & Souza, R. M. (2016). Basu–Dhar bivariate geometric distribution in the presence of covariates and censored data: a Bayesian approach. *Journal of Applied Statistics*, **43**, 9, 1636-1648.

de Oliveira, R. P., & Achcar, J. A. (2018). Basu-Dhar's bivariate geometric distribution in presence of censored data and covariates: some computational aspects. *Electronic Journal of Applied Statistical Analysis*, **11**, 1, 108-136.

de Oliveira, R. P., Achcar, J. A., Peralta, D., & Mazucheli, J. (2018). Discrete and continuous bivariate lifetime models in presence of cure rate: a comparative study under Bayesian approach. *Journal of Applied Statistics*, 1-19.

Examples

```
covbivgeo(theta = c(0.5, 0.5, 0.7))
# [1] 0.1506186
covbivgeo(theta = c(0.2, 0.5, 0.7))
# [1] 0.04039471
covbivgeo(theta = c(0.8, 0.9, 0.1))
# [1] 0.0834061
covbivgeo(theta = c(0.9, 0.9, 0.9))
# [1] 7.451626
```

dbivgeo

Joint Probability Mass Function for the Basu-Dhar Bivariate Geometric Distribution

Description

This function computes the joint probability mass function of the Basu-Dhar bivariate geometric distribution for arbitrary parameter values.

Usage

```
dbivgeo1(x, y = NULL, theta, log = FALSE)
dbivgeo2(x, y = NULL, theta, log = FALSE)
```

Arguments

x	matrix or vector containing the data. If x is a matrix then it is considered as x the first column and y the second column (y argument need be setted to NULL). Additional columns and y are ignored.
y	vector containing the data of y. It is used only if x is also a vector. Vectors x and y should be of equal length.
theta	vector (of length 3) containing values of the parameters θ_1, θ_2 and θ_3 of the Basu-Dhar bivariate Geometric distribution. The parameters are restricted to $0 < \theta_i < 1, i = 1, 2$ and $0 < \theta_3 \leq 1$.
log	logical argument for calculating the log probability or the probability function. The default value is FALSE.

Details

The joint probability mass function for a random vector (X, Y) following a Basu-Dhar bivariate geometric distribution could be written in two forms. The first form is described by:

$$P(X = x, Y = y) = \theta_1^{x-1} \theta_2^{y-1} \theta_3^{z_1} - \theta_1^x \theta_2^{y-1} \theta_3^{z_2} - \theta_1^{x-1} \theta_2^y \theta_3^{z_3} + \theta_1^x \theta_2^y \theta_3^{z_4}$$

where $x, y > 0$ are positive integers and $z_1 = \max(x - 1, y - 1), z_2 = \max(x, y - 1), z_3 = \max(x - 1, y), z_4 = \max(x, y)$. The second form is given by the conditions:

If $X < Y$, then

$$P(X = x, Y = y) = \theta_1^{x-1} (\theta_2 \theta_3)^{y-1} (1 - \theta_2 \theta_3) (1 - \theta_1)$$

If $X = Y$, then

$$P(X = x, Y = y) = (\theta_1 \theta_2 \theta_3)^{x-1} (1 - \theta_1 \theta_3 - \theta_2 \theta_3 + \theta_1 \theta_2 \theta_3)$$

If $X > Y$, then

$$P(X = x, Y = y) = \theta_2^{y-1} (\theta_1 \theta_3)^{x-1} (1 - \theta_1 \theta_3) (1 - \theta_2)$$

Value

[dbivgeo1](#) gives the values of the probability mass function using the first form of the joint pmf.

[dbivgeo2](#) gives the values of the probability mass function using the second form of the joint pmf.

Invalid arguments will return an error message.

Author(s)

Ricardo P. Oliveira <rpuziol.oliveira@gmail.com>

Jorge Alberto Achcar <achcar@fmrp.usp.br>

Source

[dbivgeo1](#) and [dbivgeo2](#) are calculated directly from the definitions.

References

- Basu, A. P., & Dhar, S. K. (1995). Bivariate geometric distribution. *Journal of Applied Statistical Science*, **2**, 1, 33-44.
- Li, J., & Dhar, S. K. (2013). Modeling with bivariate geometric distributions. *Communications in Statistics-Theory and Methods*, **42**, **2**, 252-266.
- Achcar, J. A., Davarzani, N., & Souza, R. M. (2016). Basu–Dhar bivariate geometric distribution in the presence of covariates and censored data: a Bayesian approach. *Journal of Applied Statistics*, **43**, **9**, 1636-1648.
- de Oliveira, R. P., & Achcar, J. A. (2018). Basu-Dhar's bivariate geometric distribution in presence of censored data and covariates: some computational aspects. *Electronic Journal of Applied Statistical Analysis*, **11**, **1**, 108-136.

See Also

[Geometric](#) for the univariate geometric distribution.

Examples

```
# If x and y are integer numbers:

dbivgeo1(x = 1, y = 2, theta = c(0.2, 0.4, 0.7), log = FALSE)
# [1] 0.16128
dbivgeo2(x = 1, y = 2, theta = c(0.2, 0.4, 0.7), log = FALSE)
# [1] 0.16128

# If x is a matrix:

matr <- matrix(c(1,2,3,5), ncol = 2)

dbivgeo1(x = matr, y = NULL, theta = c(0.2,0.4,0.7), log = FALSE)
# [1] 0.0451584000 0.0007080837
dbivgeo2(x = matr, y = NULL, theta = c(0.2,0.4,0.7), log = FALSE)
# [1] 0.0451584000 0.0007080837

# If log = TRUE:

dbivgeo1(x = 1, y = 2, theta = c(0.2, 0.4, 0.7), log = TRUE)
# [1] -1.824613
dbivgeo2(x = 1, y = 2, theta = c(0.2, 0.4, 0.7), log = TRUE)
# [1] -1.824613
```


Description

This function computes the joint probability mass function of the Basu-Dhar bivariate geometric distribution assuming arbitrary parameter values in presence of cure fraction.

Usage

```
dbivgeocure(x, y, theta, phi11, log = FALSE)
```

Arguments

x	matrix or vector containing the data. If x is a matrix then it is considered as x the first column and y the second column (y argument need be setted to NULL). Additional columns and y are ignored.
y	vector containing the data of y. It is used only if x is also a vector. Vectors x and y should be of equal length.
theta	vector (of length 3) containing values of the parameters θ_1, θ_2 and θ_3 of the Basu-Dhar bivariate Geometric distribution. The parameters are restricted to $0 < \theta_i < 1, i = 1, 2$ and $0 < \theta_3 \leq 1$.
phi11	real number containing the value of the cure fraction incidence parameter ϕ_{11} restricted to $0 < \phi_{11} < 1$ and $\phi_{11} + \phi_{10} + \phi_{01} + \phi_{00} = 1$ where ϕ_{10}, ϕ_{01} and ϕ_{00} are the complementary cure fraction incidence parameters for the joint cdf and sf functions.
log	logical argument for calculating the log probability or the probability function. The default value is FALSE.

Details

The joint probability mass function for a random vector (X, Y) following a Basu-Dhar bivariate geometric distribution in presence of cure fraction could be written as:

$$P(X = x, Y = y) = \phi_{11}(\theta_1^{x-1}\theta_2^{y-1}\theta_3^{z_1} - \theta_1^x\theta_2^{y-1}\theta_3^{z_2} - \theta_1^{x-1}\theta_2^y\theta_3^{z_3} + \theta_1^x\theta_2^y\theta_3^{z_4})$$

where $x, y > 0$ are positive integers and $z_1 = \max(x - 1, y - 1), z_2 = \max(x, y - 1), z_3 = \max(x - 1, y), z_4 = \max(x, y)$.

Value

[dbivgeocure](#) gives the values of the probability mass function in presence of cure fraction.

Invalid arguments will return an error message.

Author(s)

Ricardo P. Oliveira <rpuziol.oliveira@gmail.com>

Jorge Alberto Achcar <achcar@fmrp.usp.br>

Source

[dbivgeocure](#) is calculated directly from the definition.

References

Basu, A. P., & Dhar, S. K. (1995). Bivariate geometric distribution. *Journal of Applied Statistical Science*, **2**, 1, 33-44.

Achcar, J. A., Davarzani, N., & Souza, R. M. (2016). Basu–Dhar bivariate geometric distribution in the presence of covariates and censored data: a Bayesian approach. *Journal of Applied Statistics*, **43**, 9, 1636-1648.

de Oliveira, R. P., & Achcar, J. A. (2018). Basu-Dhar's bivariate geometric distribution in presence of censored data and covariates: some computational aspects. *Electronic Journal of Applied Statistical Analysis*, **11**, 1, 108-136.

de Oliveira, R. P., Achcar, J. A., Peralta, D., & Mazucheli, J. (2018). Discrete and continuous bivariate lifetime models in presence of cure rate: a comparative study under Bayesian approach. *Journal of Applied Statistics*, 1-19.

See Also

[Geometric](#) for the univariate geometric distribution.

Examples

```
# If log = FALSE:

dbivgeocure(x = 1, y = 2, theta = c(0.2, 0.4, 0.7), phi11 = 0.4, log = FALSE)
# [1] 0.064512

# If log = TRUE:

dbivgeocure(x = 1, y = 2, theta = c(0.2, 0.4, 0.7), phi11 = 0.4, log = TRUE)
# [1] -2.740904
```

mombivgeo

Moments Estimator for the Basu-Dhar Bivariate Geometric Distribution

Description

This function computes the estimators based on the method of the moments for each parameter of the Basu-Dhar bivariate geometric distribution.

Usage

```
mombivgeo(x, y)
```

Arguments

- x matrix or vector containing the data. If x is a matrix then it is considered as x the first column and y the second column (y argument need be setted to NULL). Additional columns and y are ignored.
- y vector containing the data of y. It is used only if x is also a vector. Vectors x and y should be of equal length.

Details

The moments estimators of $\theta_1, \theta_2, \theta_3$ of the Basu-Dhar bivariate geometric distribution are given by:

$$\hat{\theta}_1 = \frac{\bar{Y}(1 - \bar{W})}{\bar{W}(1 - \bar{Y})}$$

$$\hat{\theta}_2 = \frac{\bar{X}(\bar{W} - 1)}{\bar{W}(\bar{X} - 1)}$$

$$\hat{\theta}_3 = \frac{\bar{X}(\bar{X} - 1)(\bar{Y} - 1)}{(\bar{W} - 1)\bar{X}\bar{Y}}$$

Value

[mombivgeo](#) gives the values of the moments estimator.

Invalid arguments will return an error message.

Author(s)

Ricardo P. Oliveira <rpuziol.oliveira@gmail.com>

Jorge Alberto Achcar <achcar@fmrp.usp.br>

Source

[mombivgeo](#) is calculated directly from the definition.

References

Basu, A. P., & Dhar, S. K. (1995). Bivariate geometric distribution. *Journal of Applied Statistical Science*, **2**, 1, 33-44.

Li, J., & Dhar, S. K. (2013). Modeling with bivariate geometric distributions. *Communications in Statistics-Theory and Methods*, **42**, **2**, 252-266.

Achcar, J. A., Davarzani, N., & Souza, R. M. (2016). Basu–Dhar bivariate geometric distribution in the presence of covariates and censored data: a Bayesian approach. *Journal of Applied Statistics*, **43**, **9**, 1636-1648.

de Oliveira, R. P., & Achcar, J. A. (2018). Basu-Dhar's bivariate geometric distribution in presence of censored data and covariates: some computational aspects. *Electronic Journal of Applied Statistical Analysis*, **11**, **1**, 108-136.

See Also

[Geometric](#) for the univariate geometric distribution.

Examples

```
# Generate the data set:

set.seed(123)
x1 <- rbivgeo1(n = 1000, theta = c(0.5, 0.5, 0.7))
set.seed(123)
x2 <- rbivgeo2(n = 1000, theta = c(0.5, 0.5, 0.7))

# Compute de moment estimator by:

mombivgeo(x = x1, y = NULL) # For data set x1
#           [,1]
# theta1 0.5053127
# theta2 0.5151873
# theta3 0.6640406

mombivgeo(x = x2, y = NULL) # For data set x2
#           [,1]
# theta1 0.4922327
# theta2 0.5001577
# theta3 0.6993893
```

pbivgeo

Joint Cumulative Function for the Basu-Dhar Bivariate Geometric Distribution

Description

This function computes the joint cumulative function of the Basu-Dhar bivariate geometric distribution assuming arbitrary parameter values.

Usage

```
pbivgeo(x, y, theta, lower.tail = TRUE)
```

Arguments

x	matrix or vector containing the data. If x is a matrix then it is considered as x the first column and y the second column (y argument need be setted to NULL). Additional columns and y are ignored.
y	vector containing the data of y. It is used only if x is also a vector. Vectors x and y should be of equal length.
theta	vector (of length 3) containing values of the parameters θ_1, θ_2 and θ_3 of the Basu-Dhar bivariate Geometric distribution. The parameters are restricted to $0 < \theta_i < 1, i = 1, 2$ and $0 < \theta_3 \leq 1$.

lower.tail logical; If TRUE (default), probabilities are $P(X \leq x, Y \leq y)$ otherwise $P(X > x, Y > y)$.

Details

The joint cumulative function for a random vector (X, Y) following a Basu-Dhar bivariate geometric distribution could be written as:

$$P(X \leq x, Y \leq y) = 1 - (\theta_1 \theta_3)^x - (\theta_2 \theta_3)^y + \theta_1^x \theta_2^y \theta_3^{\max(x,y)}$$

and the joint survival function is given by:

$$P(X > x, Y > y) = \theta_1^x \theta_2^y \theta_3^{\max(x,y)}$$

Value

[pbivgeo](#) gives the values of the cumulative function.

Invalid arguments will return an error message.

Author(s)

Ricardo P. Oliveira <rpuziol.oliveira@gmail.com>

Jorge Alberto Achcar <achcar@fmrp.usp.br>

Source

[pbivgeo](#) is calculated directly from the definition.

References

Basu, A. P., & Dhar, S. K. (1995). Bivariate geometric distribution. *Journal of Applied Statistical Science*, **2**, 1, 33-44.

Li, J., & Dhar, S. K. (2013). Modeling with bivariate geometric distributions. *Communications in Statistics-Theory and Methods*, **42**, **2**, 252-266.

Achcar, J. A., Davarzani, N., & Souza, R. M. (2016). Basu–Dhar bivariate geometric distribution in the presence of covariates and censored data: a Bayesian approach. *Journal of Applied Statistics*, **43**, 9, 1636-1648.

de Oliveira, R. P., & Achcar, J. A. (2018). Basu-Dhar's bivariate geometric distribution in presence of censored data and covariates: some computational aspects. *Electronic Journal of Applied Statistical Analysis*, **11**, 1, 108-136.

See Also

[Geometric](#) for the univariate geometric distribution.

Examples

```
# If x and y are integer numbers:

pbivgeo(x = 1, y = 2, theta = c(0.2, 0.4, 0.7), lower.tail = TRUE)
# [1] 0.79728

# If x is a matrix:

matr <- matrix(c(1,2,3,5), ncol = 2)
pbivgeo(x = matr, y = NULL, theta = c(0.2,0.4,0.7), lower.tail = TRUE)
# [1] 0.8424384 0.9787478

# If lower.tail = FALSE:

pbivgeo(x = 1, y = 2, theta = c(0.2, 0.4, 0.7), lower.tail = FALSE)
# [1] 0.01568

matr <- matrix(c(1,2,3,5), ncol = 2)
pbivgeo(x = matr, y = NULL, theta = c(0.9,0.4,0.7), lower.tail = FALSE)
# [1] 0.01975680 0.00139404
```

pbivgeocure

Joint Cumulative Function for the Basu-Dhar Bivariate Geometric Distribution in Presence of Cure Fraction

Description

This function computes the joint cumulative function of the Basu-Dhar bivariate geometric distribution assuming arbitrary parameter values in presence of cure fraction.

Usage

```
pbivgeocure(x, y, theta, phi, lower.tail = TRUE)
```

Arguments

x	matrix or vector containing the data. If x is a matrix then it is considered as x the first column and y the second column (y argument need be setted to NULL). Additional columns and y are ignored.
y	vector containing the data of y. It is used only if x is also a vector. Vectors x and y should be of equal length.
theta	vector (of length 3) containing values of the parameters θ_1, θ_2 and θ_3 of the Basu-Dhar bivariate Geometric distribution. The parameters are restricted to $0 < \theta_i < 1, i = 1, 2$ and $0 < \theta_3 \leq 1$.
phi	vector (of length 4) containing values of the cure fraction incidence parameters $\phi_{11}, \phi_{10}, \phi_{01}$ and ϕ_{00} . The parameters are restricted to $\phi_{11} + \phi_{10} + \phi_{01} + \phi_{00} = 1$.

`lower.tail` logical; If TRUE (default), probabilities are $P(X \leq x, Y \leq y)$ otherwise $P(X > x, Y > y)$.

Details

The joint cumulative function for a random vector (X, Y) following a Basu-Dhar bivariate geometric distribution in presence of cure fraction could be written as:

$$P(X \leq x, Y \leq y) = 1 - (\phi_{11} + \phi_{10})(\theta_1\theta_3)^x - (\phi_{01} + \phi_{00}) - (\phi_{11} + \phi_{01})(\theta_2\theta_3)^y - (\phi_{10} + \phi_{00}) \\ + \phi_{11}(\theta_1^x\theta_2^y\theta_3^{\max(x,y)}) + \phi_{10}(\theta_1\theta_3)^x + \phi_{01}(\theta_2\theta_3)^y + \phi_{00}$$

and the joint survival function is given by:

$$P(X > x, Y > y) = \phi_{11}(\theta_1^x\theta_2^y\theta_3^{\max(x,y)}) + \phi_{10}(\theta_1\theta_3)^x + \phi_{01}(\theta_2\theta_3)^y + \phi_{00}$$

Value

[pbivgeocure](#) gives the values of the cumulative function in presence of cure fraction.

Invalid arguments will return an error message.

Author(s)

Ricardo P. Oliveira <rpuziol.oliveira@gmail.com>

Jorge Alberto Achcar <achcar@fmrp.usp.br>

Source

[pbivgeocure](#) is calculated directly from the definition.

References

Basu, A. P., & Dhar, S. K. (1995). Bivariate geometric distribution. *Journal of Applied Statistical Science*, **2**, 1, 33-44.

Achcar, J. A., Davarzani, N., & Souza, R. M. (2016). Basu–Dhar bivariate geometric distribution in the presence of covariates and censored data: a Bayesian approach. *Journal of Applied Statistics*, **43**, 9, 1636-1648.

de Oliveira, R. P., & Achcar, J. A. (2018). Basu-Dhar's bivariate geometric distribution in presence of censored data and covariates: some computational aspects. *Electronic Journal of Applied Statistical Analysis*, **11**, 1, 108-136.

de Oliveira, R. P., Achcar, J. A., Peralta, D., & Mazucheli, J. (2018). Discrete and continuous bivariate lifetime models in presence of cure rate: a comparative study under Bayesian approach. *Journal of Applied Statistics*, 1-19.

See Also

[Geometric](#) for the univariate geometric distribution.

Examples

```
# If lower.tail = TRUE:

pbivgeocure(x = 1, y = 2, theta = c(0.2, 0.4, 0.7), phi = c(0.2, 0.3, 0.3, 0.2), lower.tail = TRUE)
# [1] 0.159456

matr <- matrix(c(1,2,3,5), ncol = 2)
pbivgeocure(x=matr,y=NULL,theta=c(0.2, 0.4, 0.7),phi=c(0.2, 0.3, 0.3, 0.2),lower.tail = TRUE)
# [1] 0.1684877 0.1957496

# If lower.tail = FALSE:

pbivgeocure(x = 1, y = 2, theta = c(0.2, 0.4, 0.7), phi = c(0.2, 0.3, 0.3, 0.2), lower.tail = FALSE)
# [1] 0.268656

matr <- matrix(c(1,2,3,5), ncol = 2)
pbivgeocure(x=matr,y=NULL,theta=c(0.2, 0.4, 0.7),phi=c(0.2, 0.3, 0.3, 0.2),lower.tail = FALSE)
# [1] 0.2494637 0.2064101
```

rbivgeo

Generates Random Deviates from the Basu-Dhar Bivariate Geometric Distribution

Description

This function generates random values from the Basu-Dhar bivariate geometric distribution assuming arbitrary parameter values.

Usage

```
rbivgeo1(n, theta)
rbivgeo2(n, theta)
```

Arguments

n	number of observations. If $\text{length}(n) > 1$, the length is taken to be the number required.
theta	vector (of length 3) containing values of the parameters θ_1, θ_2 and θ_3 of the Basu-Dhar bivariate Geometric distribution. The parameters are restricted to $0 < \theta_i < 1, i = 1, 2$ and $0 < \theta_3 \leq 1$.

Details

The conditional distribution of X given Y is given by:

If $X < Y$, then

$$P(X = x|Y = y) = \theta_1^{x-1}(1 - \theta_1)$$

If $X = Y$, then

$$P(X = x|Y = y) = \frac{\theta_1^{x-1}(1 - \theta_1\theta_3 - \theta_2\theta_3 + \theta_1\theta_2\theta_3)}{1 - \theta_2\theta_3}$$

If $X > Y$, then

$$P(X = x|Y = y) = \frac{\theta_1^{x-1}\theta_3^{x-y}(1 - \theta_1\theta_3)(1 - \theta_2)}{1 - \theta_2\theta_3}$$

Value

`rbivgeo1` and `rbivgeo2` generate random deviates from the Bash-Dhar bivariate geometric distribution. The length of the result is determined by `n`, and is the maximum of the lengths of the numerical arguments for the other functions.

Invalid arguments will return an error message.

Author(s)

Ricardo P. Oliveira <rpuziol.oliveira@gmail.com>

Jorge Alberto Achcar <achcar@fmrp.usp.br>

Source

`rbivgeo1` generates random deviates using the inverse transformation method. Returns a matrix that the first column corresponds to X generated random values and the second column corresponds to Y generated random values.

`rbivgeo2` generates random deviates using the shock model. Returns a matrix that the first column corresponds to X generated random values and the second column corresponds to Y generated random values. See Marshall and Olkin (1967) for more details.

References

- Marshall, A. W., & Olkin, I. (1967). A multivariate exponential distribution. *Journal of the American Statistical Association*, **62**, 317, 30-44.
- Basu, A. P., & Dhar, S. K. (1995). Bivariate geometric distribution. *Journal of Applied Statistical Science*, **2**, 1, 33-44.
- Li, J., & Dhar, S. K. (2013). Modeling with bivariate geometric distributions. *Communications in Statistics-Theory and Methods*, **42**, 2, 252-266.
- Achcar, J. A., Davarzani, N., & Souza, R. M. (2016). Basu–Dhar bivariate geometric distribution in the presence of covariates and censored data: a Bayesian approach. *Journal of Applied Statistics*, **43**, 9, 1636-1648.
- de Oliveira, R. P., & Achcar, J. A. (2018). Basu-Dhar's bivariate geometric distribution in presence of censored data and covariates: some computational aspects. *Electronic Journal of Applied Statistical Analysis*, **11**, 1, 108-136.

See Also

[Geometric](#) for the univariate geometric distribution.

Examples

```
rbivgeo1(n = 10, theta = c(0.5, 0.5, 0.7))
#      [,1] [,2]
# [1,]  2   1
# [2,]  3   1
# [3,]  1   1
# [4,]  1   1
# [5,]  2   2
# [6,]  1   3
# [7,]  2   2
# [8,]  1   1
# [9,]  1   1
# [10,] 2   2
```

```
rbivgeo2(n = 10, theta = c(0.5, 0.5, 0.7))
#      [,1] [,2]
# [1,]  1   1
# [2,]  2   1
# [3,]  2   1
# [4,]  4   1
# [5,]  1   1
# [6,]  2   2
# [7,]  3   2
# [8,]  3   1
# [9,]  3   2
# [10,] 1   1
```

Index

cfbivgeo, [2](#), [2](#), [3](#)
corbivgeo, [3](#), [4](#)
covbivgeo, [5](#), [5](#), [6](#)

dbivgeo, [6](#)
dbivgeo1, [7](#)
dbivgeo1 (dbivgeo), [6](#)
dbivgeo2, [7](#)
dbivgeo2 (dbivgeo), [6](#)
dbivgeocure, [8](#), [9](#)

Geometric, [8](#), [10](#), [12](#), [13](#), [15](#), [17](#)

mombivgeo, [10](#), [11](#)

pbivgeo, [12](#), [13](#)
pbivgeocure, [14](#), [15](#)

rbivgeo, [16](#)
rbivgeo1, [17](#)
rbivgeo1 (rbivgeo), [16](#)
rbivgeo2, [17](#)
rbivgeo2 (rbivgeo), [16](#)